

On Sobolev type orthogonal polynomials  
with unbounded support: asymptotic properties

M. Alfaro<sup>a</sup>, J. J. Moreno-Balcázar<sup>b</sup>, A. Peña<sup>a</sup> and M. L. Rezola<sup>a</sup>

<sup>a</sup>Depto. de Matemáticas and IUMA. Universidad de Zaragoza (Spain).

<sup>b</sup>Depto. de Estadística y Matemática Aplicada and ICI. Universidad de Almería (Spain).

**Abstract**

In this expository paper we present a survey about asymptotic properties of Sobolev type orthogonal polynomials with unbounded support.

Key words: Sobolev orthogonal polynomials, Relative asymptotics; Mehler–Heine type formulas; zeros; Bessel functions.

2000MSC: 42C05, 33C45.

**1 Introduction**

The theory of orthogonal polynomials is a very interesting field in mathematics with important applications to numerical analysis, physics, probability, and statistics among other ones. Orthogonal polynomials are connected with topics like moment problems, mechanical quadratures, continued fractions, spectral methods, quantum mechanics and many other concepts.

Usually, in this theory, the orthogonality is considered with respect to a positive linear functional defined on the linear space of polynomials or, according to the Riesz representation theorem, with respect to a positive measure. Let  $\mu$  be a finite positive Borel measure supported on an interval  $I$  in the real line, we say that the sequence of polynomials  $\{P_n\}_{n \geq 0}$  is a sequence of orthogonal polynomials (o.p.) with respect to either the measure  $\mu$  or the inner product  $(f, g) = \int_I f g d\mu$  if, for all  $n \geq 0$ ,  $\deg P_n = n$  and

(i)  $(P_n, P_m) = 0, n \neq m,$

(ii)  $(P_n, P_n) > 0, n \geq 0.$

Along the paper such a kind of inner products will be called standard inner products. They have the following remarkable property:  $(xp, q) = (p, xq)$ , for all polynomials  $p, q$ . As a consequence, the corresponding standard orthogonal polynomials have nice properties such as the three-term recurrence relation, the summation formula, the interlacing properties of the zeros, etc. From a numerical point of view, a useful consequence is that a Gaussian mechanical quadrature formula has exact precision when we take as nodes the zeros of appropriate standard o.p..

Nonstandard inner products have also been considered in the literature. In particular, the so-called Sobolev inner products that are of the form

$$(f, g) = \int f g d\mu_0 + \sum_{i=1}^r \int f^{(i)} g^{(i)} d\mu_i,$$

where  $\{\mu_i\}_{i=0}^r$  are finite positive Borel measures supported on the real line and the functions  $f$  and  $g$  belong to the Sobolev space:

$$W^{2,r}(\mu_0, \mu_1, \dots, \mu_r) := \{f : \int |f|^2 d\mu_0 + \sum_{i=1}^r \int |f^{(i)}|^2 d\mu_i < +\infty\}.$$

Studied by the first time in the forties of the last century, the Sobolev orthogonal polynomials have been object of an increasing interest, approximately, in the last 20 years. Obviously, Sobolev inner products are nonstandard and therefore Sobolev o.p. loose the “good” properties of the standard o.p. However, it is interesting to study these “strange” polynomials that supply us with situations different from the standard ones: no three-term recurrence relation, zeros out of the convex hull of the support of the orthogonality measure including, some times, complex zeros, and so one.

Furthermore, some applications of the Sobolev orthogonality in the theory of standard o.p. are known, for instance, classical polynomials (Jacobi or Laguerre polynomials) with nonclassical parameters are not orthogonal in the usual sense but they are orthogonal with respect to Sobolev inner products (see among others [1] or [17]) and also, Sobolev o.p. in two real variables are solutions of some partial differential equations (see [9], [14], [19] or [24]).

In this paper we are concerned with the so-called Sobolev type (or discrete Sobolev) orthogonal polynomials, that is, polynomials orthogonal with respect to a Sobolev inner product in which  $\{\mu_i\}_{i=1}^r$  are Dirac’s deltas or, in general, discrete measures. More concretely, we consider an inner product of the form

$$(f, g) = \int f(x)g(x)d\mu(x) + \sum_{i=0}^r M_i f^{(i)}(c)g^{(i)}(c),$$

where  $\mu$  is a finite positive Borel measure,  $c \in \mathbb{R}$  and  $M_i \geq 0$  for  $i = 0, 1, \dots, r$ . In the sequel, we denote by  $\{Q_{n,r}\}_{n \geq 0}$  the corresponding sequence of o.p. with the same leading coefficient as the standard o.p. with respect to  $\mu$ .

More general products where cross-product terms appear in the discrete part (the non-diagonal case) have also been studied. But, recently in [18] the authors prove that every symmetric bilinear form can be reduced to a diagonal case, that is, an inner product without cross-product terms.

In some sense, Sobolev type o.p. are not so far than the standard o.p. since there is the possibility to transform the Sobolev type orthogonality into the standard quasi-orthogonality. As a consequence, several properties of the standard o.p. are partially recovered for the Sobolev type o.p.: they satisfy a  $2r + 3$  term recurrence relation (see, [13]) and have partial interlacing properties of the zeros ([2]).

Since the polynomial  $Q_{n,r}$  is quasi-orthogonal of order  $r+1$  with respect to the measure  $\mu$  it can be expressed as a linear combination (with a fixed number of terms:  $r + 2$ ) of standard orthogonal polynomials  $R_n$  corresponding to the modified measure  $(x - c)^{r+1}d\mu$ , that is,

$$Q_{n,r}(x) = \sum_{j=0}^{r+1} a_n^j R_{n-j}(x). \quad (1)$$

One of the topics in the theory is to compare the Sobolev type o.p. with the standard o.p. (with respect to  $\mu$ ) to investigate how the addition of the discrete part in the inner product influences the orthogonal system. Many formal results are known for the polynomials  $Q_{n,r}$ : recurrence relation, differential formulas, location of zeros, and so on. However, little is known about the asymptotic properties and most of the general results have been obtained when the support of  $\mu$  ( $\text{supp } \mu$ ) is a bounded set. For instance, in [20], the authors assume that  $\mu$  is a measure of bounded support for which the asymptotic behaviour of the corresponding o.p. is known; the most relevant class of this type is the Nevai class  $M(0, 1)$  of o.p. with appropriately converging recurrence coefficients. There, the relative asymptotics is studied when the mass point  $c \notin \text{supp } \mu$ . The case  $c \in \text{supp } \mu$  has been considered in [22].

What happens is that in the bounded case, all the connexion coefficients  $a_n^j$  in (1) are bounded and the orthogonal polynomials  $R_n$  have an adequate finite ratio asymptotics: these two facts allow us to study each term of (1) separately, in order to get the relative asymptotics for  $Q_{n,r}$  (see [20] and [22] where this technique is developed). However, the situation is quite different if we deal with the unbounded case because when we try to obtain the relative asymptotics with the techniques used for the bounded case and we take into account the ratio asymptotics for the polynomials  $R_n$ , we come across a serious problem. Indeed, we find that the idea that each term of (1) has a finite limit could not work now, in fact, as we will see later, for the Laguerre-Sobolev type o.p. each term of (1) tends to infinity, all of them being the same order, but with an alternating sign.

The aim of this paper is to describe the current state of the asymptotic properties for Sobolev type polynomials when  $\text{supp } \mu$  is unbounded. Mainly we will analyze the case

when  $\mu$  is the Laguerre probability measure ( $d\mu(x) = \frac{x^\alpha e^{-x}}{\Gamma(\alpha+1)} dx$  with  $\alpha > -1$ ) and  $c = 0$ , that is, the Laguerre–Sobolev o.p. This choice for  $c$  is due to the fact that the point  $x = 0$  is a singularity of the differential equation satisfied by the classical Laguerre polynomials.

Therefore, we will deal with classical Laguerre polynomials, that is, polynomials orthogonal with respect to the inner product

$$(p, q) = \frac{1}{\Gamma(\alpha + 1)} \int_0^\infty p(x)q(x) x^\alpha e^{-x} dx, \quad \alpha > -1.$$

in the space of all polynomials with real coefficients. We will denote by  $L_n^\alpha$  the  $n$ th Laguerre polynomial with  $(-1)^n/n!$  as leading coefficient. Although many of the properties of Laguerre polynomials can be seen, for example, in the books of Chihara [10] and Szegő [23], we remind that the classical Laguerre polynomials with the normalization above quoted are defined by

$$L_n^\alpha(x) = \sum_{k=0}^n \binom{n+\alpha}{n-k} \frac{(-1)^k x^k}{k!},$$

and their derivatives satisfy

$$(L_n^\alpha)^{(k)}(x) = (-1)^k L_{n-k}^{\alpha+k}(x). \quad (2)$$

The evaluations at  $x = 0$  of the polynomial  $L_n^\alpha$  and its successive derivatives are given by

$$(L_n^\alpha)^{(k)}(0) = \frac{(-1)^k n!}{(n-k)!} \frac{\Gamma(\alpha+1)}{\Gamma(\alpha+k+1)} L_n^\alpha(0) = \frac{(-1)^k}{(n-k)!} \frac{\Gamma(n+\alpha+1)}{\Gamma(\alpha+k+1)}. \quad (3)$$

From Perron's formula in Szegő's book [23], the following asymptotic results can be deduced:

$$\frac{L_{n-1}^\alpha(x)}{L_n^\alpha(x)} \Rightarrow 1, \quad x \in \mathbb{C} \setminus [0, \infty), \quad (4)$$

$$\frac{n^{1/2} L_n^\alpha(x)}{L_n^{\alpha+1}(x)} \Rightarrow \sqrt{-x}, \quad x \in \mathbb{C} \setminus [0, \infty). \quad (5)$$

where the symbol  $f_n(x) \Rightarrow f(x)$ ,  $x \in A$ , denotes that the sequence  $\{f_n\}$  converges to  $f$  uniformly on compact subsets of  $A$ .

Later on we will use the symbol  $f(x) \sim g(x)$  ( $x \rightarrow a$ ) if  $\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = 1$ .

## 2 Laguerre–Sobolev type polynomials

From now on  $\{Q_{n,r}\}_{n \geq 0}$  denotes the sequence of polynomials orthogonal with respect to an inner product of the form

$$(p, q)_r = \frac{1}{\Gamma(\alpha + 1)} \int_0^\infty p(x)q(x) x^\alpha e^{-x} dx + \sum_{i=0}^r M_i p^{(i)}(0) q^{(i)}(0), \quad (6)$$

where  $\alpha > -1$  and  $M_i > 0$ ,  $i = 0, \dots, r$ , with leading coefficient  $(-1)^n/n!$ . Notice that all the masses in the discrete part are positive.

Observe that, in fact,  $(\cdot, \cdot)_r$  and  $Q_{n,r}$  also depend on the parameter  $\alpha$  but for simplicity we have omitted it in the notations.

These families of o.p. were considered by the first time by Koekoek and Meijer (see, among others, [15] and [16]) although no asymptotics were studied. The first asymptotic results for Laguerre–Sobolev type o.p. appear in [8]: exterior asymptotics, asymptotics on compact subsets of  $(0, +\infty)$ , exterior Plancherel–Rotach type asymptotics, Mehler–Heine type formulas and convergence of their zeros are obtained, but only for  $r = 0$  and  $r = 1$ . Concerning the Mehler–Heine type formulas, with  $r = 1$  and  $M_0, M_1 > 0$  the authors found a behaviour pattern and they established a conjecture. A survey including these results can be seen in [21]. Some of these properties were proved for the non–diagonal case with  $r = 1$  in [6] and [7], and later on in [11].

In all these papers the basic tool was the algebraic expression

$$Q_{n,1}(x) = B_0(n)L_n^\alpha(x) + B_1(n)xL_{n-1}^{\alpha+2}(x) + B_2(n)x^2L_{n-2}^{\alpha+4}(x)$$

where the coefficients  $B_i(n)$  were given explicitly in [16].

In a discrete Laguerre–Sobolev inner product with an arbitrary number of terms, the problem is that we only have an algebraic expression given in [15], but not the explicit expression of the coefficients  $B_i(n)$ , of which we only know that they are a non trivial solution of a system with  $r + 1$  equations and  $r + 2$  unknowns.

Asymptotic properties of Sobolev orthogonal polynomials with respect to a general inner product as (6), that is, with an arbitrary number of masses, have been studied in [5] where, in particular, the conjecture established in [8] is proved to be true. In the sequel we summarize the results obtained there.

As we have already said, the interest lies in knowing the differences in the asymptotic behaviour between the Laguerre polynomials and the Sobolev polynomials  $Q_{n,r}$ . Intuitively one can imagine that these differences in the complex plane should be around the perturbation of the standard inner product involved in the Sobolev inner product, that is, around the origin and therefore we cannot expect that the addition of a finite number of masses to the inner product produces a modification in the global behaviour of the polynomials. A result which supports this intuition is Lemma 2 in [5] where it has been proved:

**Lemma 1** *Let  $Q_{n,r}$  be the polynomials orthogonal with respect to (6) with leading coefficients  $(-1)^n/n!$ . Then the following statements hold:*

(a) For  $0 \leq k \leq r$ ,

$$Q_{n,r}^{(k)}(0) \sim \frac{C_{r,k}}{n^{\alpha+2k+1}}(L_n^\alpha)^{(k)}(0),$$

where  $C_{r,k}$  is a nonzero real number independent of  $n$ .

(b) For  $k \geq r + 1$ ,

$$Q_{n,r}^{(k)}(0) \sim \frac{k!}{(k - (r + 1))!} \frac{\Gamma(\alpha + k + 1)}{\Gamma(\alpha + r + k + 2)} (L_n^\alpha)^{(k)}(0).$$

(c)

$$(Q_{n,r}, Q_{n,r})_r \sim \|L_n^\alpha\|^2.$$

**Remark 1.** Observe that both Laguerre and Laguerre–Sobolev type polynomials have asymptotically the same global size (from the point of view of the norm), while the size of the successive derivatives at the point  $x = 0$  is affected by the discrete part of the inner product but only whenever the order of the derivatives corresponds to a positive mass.

Now we analyze two other asymptotics of the polynomials  $Q_{n,r}$ : the *relative asymptotics*, which assures that both families  $Q_{n,r}$  and  $L_n^\alpha$  are identical asymptotically on compact subsets of  $\mathbb{C} \setminus [0, \infty)$ , and the so-called Mehler–Heine type formula which shows how the presence of the masses in the inner product changes the asymptotic behaviour around the origin.

As we have mentioned before, for a discrete Sobolev inner product when  $\text{supp } \mu$  is bounded, a tool to obtain some results is the relation between the Sobolev orthogonality and the standard quasi-orthogonality.

Now, in our particular case, the sequence  $\{Q_{n,r}\}_{n \geq 0}$  is quasi-orthogonal of order  $r + 1$  with respect to the Laguerre weight  $x^{\alpha+r+1}e^{-x}$ , that is,

$$\int_0^{+\infty} p(x)Q_{n,r}(x)x^{\alpha+r+1}e^{-x}dx = 0,$$

for every polynomial  $p$  with  $\deg p \leq n - (r + 1) - 1$ . Therefore, we have a *connexion formula* of the form

$$Q_{n,r}(x) = \sum_{j=0}^{r+1} a_{n,r}^j L_{n-j}^{\alpha+r+1}(x), \quad a_{n,r}^0 = 1. \quad (7)$$

In order to deduce properties of  $Q_{n,r}$  it is convenient to know the size of the *connexion coefficients*  $a_{n,r}^j$ . In [5], it has been introduced a fruitful and new technique which leads to determine their asymptotic behaviour.

Using (7), it can be obtained a new algebraic expression which relates  $\frac{Q_{n,r}^{(k+1)}(0)}{(L_n^\alpha)^{(k+1)}(0)}$  to  $\frac{Q_{n,r}^{(k)}(0)}{(L_n^\alpha)^{(k)}(0)}$  (see [5, Lemma 3]) and allows to prove:

**Theorem 1** *Let  $a_{n,r}^j$  be the connexion coefficients which appear in (7). Then, we have*

$$\lim_n a_{n,r}^j = (-1)^j \binom{r+1}{j}, \quad 0 \leq j \leq r+1. \quad (8)$$

As a token of the interest of this result we use it to deduce an asymptotics of the Laguerre–Sobolev polynomials on compact subsets of  $(0, +\infty)$ .

**Proposition 1** *The sequence  $\{n^{-(2\alpha+2r+1)/4}Q_{n,r}\}_{n\geq 1}$  is uniformly bounded on compact subsets of  $(0, +\infty)$ .*

**Proof.** The sequence  $\{n^{-\alpha/2+1/4}L_n^\alpha\}_{n\geq 1}$  is uniformly bounded on compact subsets of  $(0, +\infty)$  (see Theorem 8.22.1 in [23]), and then, for all  $j = 0, 1, \dots, r+1$ , the sequences  $\{n^{-(2\alpha+2r+1)/4}L_{n-j}^{\alpha+r+1}\}_{n\geq 1}$  are uniformly bounded on compact subsets of  $(0, +\infty)$ . From (8) and the connexion formula the result follows.  $\square$

However, it is worth noticing that the knowledge of the asymptotic behaviour of the connexion coefficients is not enough to deduce other asymptotic properties. Indeed, concerning the relative asymptotics, from (7) we have

$$\frac{Q_{n,r}(x)}{L_n^\alpha(x)} = \sum_{j=0}^{r+1} a_{n,r}^j \frac{L_{n-j}^{\alpha+r+1}(x)}{L_n^\alpha(x)}.$$

Applying Theorem 1, and (4) and (5) each term in the above sum tends to infinity with the same order but with an alternating sign, that is,

$$a_{n,r}^j \frac{L_{n-j}^{\alpha+r+1}(x)}{L_n^\alpha(x)} \sim (-1)^j \binom{r+1}{j} \left(\frac{1}{\sqrt{-x}}\right)^{r+1} n^{\frac{r+1}{2}},$$

uniformly on compact subsets of  $\mathbb{C} \setminus [0, \infty)$ .

Since the techniques used in the bounded case do not work when  $\text{supp } \mu$  is an unbounded set we proceed in a different way to prove:

**Theorem 2** *Let  $\{Q_{n,r}\}_{n\geq 0}$  be the sequence of polynomials orthogonal with respect to the inner product (6) with  $(-1)^n/n!$  as leading coefficient. Then, for  $k \geq 0$ ,*

$$\lim_n \frac{Q_{n,r}^{(k)}(x)}{(L_n^\alpha)^{(k)}(x)} = 1,$$

uniformly on compact subsets of  $\mathbb{C} \setminus [0, \infty)$ .

**Proof.** From the Fourier expansion of the polynomial  $Q_{n0}$  in terms of Laguerre polynomials, using Lemma 1, (3) and (4) the result follows for  $k = 0$ . (For more details see Theorem 1 in [5]).

The functions  $Q_{n,r}/L_n^\alpha$  are analytic in  $\mathbb{C} \setminus [0, \infty)$  and  $\frac{Q_{n,r}(x)}{L_n^\alpha(x)} \rightrightarrows 1, x \in \mathbb{C} \setminus [0, \infty)$ , then  $\left(\frac{Q_{n,r}}{L_n^\alpha}\right)'(x) \rightrightarrows 0, x \in \mathbb{C} \setminus [0, \infty)$ . Therefore,

$$\left(\frac{Q'_{n,r}(x)}{(L_n^\alpha)'(x)} - \frac{Q_{n,r}(x)}{L_n^\alpha(x)}\right) \frac{(L_n^\alpha)'(x)}{L_n^\alpha(x)} \rightrightarrows 0, x \in \mathbb{C} \setminus [0, \infty).$$

From (2), (4) and (5), we get  $\frac{(L_n^\alpha)'(x)}{L_n^\alpha(x)} \Rightarrow \infty$  and then

$$\lim_n \frac{Q'_{n,r}(x)}{(L_n^\alpha)'(x)} = \lim_n \frac{Q_{n,r}(x)}{L_n^\alpha(x)} = 1$$

uniformly on compact subsets of  $\mathbb{C} \setminus [0, \infty)$ . So, the result holds for  $k = 1$ .

Using this technique, by an induction procedure, the result follows for all  $k \geq 0$ .  $\square$

Once we know that both sequences of orthogonal polynomials,  $\{Q_{n,r}\}_{n \geq 0}$  and  $\{L_n^\alpha\}_{n \geq 0}$ , are asymptotically identical on compact subsets of  $\mathbb{C} \setminus [0, \infty)$ , we establish their differences.

To do this, we consider Mehler–Heine type formulas because they are nice tools to describe the polynomials around the origin. These kind of formulas are interesting twofold: they provide the scaled asymptotics for  $Q_{n,r}$  on compact sets of the complex plane and they supply us with asymptotic information about the location of the zeros of these polynomials in terms of the zeros of other known special functions. More precisely, applying Hurwitz’s Theorem in a straightforward way, the existence of an acceleration of the convergence of  $r + 1$  zeros of these Sobolev polynomials towards the origin can be proved.

First of all, we recall the corresponding formula for the classical Laguerre polynomials, (see [23, Th.8.1.3]):

$$n^{-\alpha} L_n^\alpha \left( \frac{x}{n} \right) \Rightarrow x^{-\alpha/2} J_\alpha(2\sqrt{x}), \quad x \in \mathbb{C}, \quad (9)$$

where  $J_\alpha$  is the Bessel function of the first kind of order  $\alpha$  ( $\alpha > -1$ ).

As it occurs in the study of the relative asymptotics, the Mehler–Heine type formulas cannot be deduced as a consequence of the connexion formula. Indeed, from (7) we have

$$n^{-\alpha} Q_{n,r} \left( \frac{x}{n} \right) = \sum_{i=0}^{r+1} a_{n,r}^i n^{-\alpha} L_{n-i}^{\alpha+r+1} \left( \frac{x}{n} \right).$$

and, applying Theorem 1 and (9), we have that each term tends to infinity with the same order but with an alternating sign.

Thus, to get the result for  $\{Q_{n,r}\}_{n \geq 0}$ , the problem should be focused on in a different way. An approach consists in to write the Taylor expansion of the polynomial  $Q_{n,r}$

$$n^{-\alpha} Q_{n,r} \left( \frac{x}{n} \right) = \sum_{k=0}^n \frac{Q_{n,r}^{(k)}(0)}{(L_n^\alpha)^{(k)}(0)} \frac{(L_n^\alpha)^{(k)}(0)}{k!} \frac{x^k}{n^{\alpha+k}},$$

and to calculate the limit applying the Lebesgue’s dominated convergence theorem. So, we need to prove that the ratios  $Q_{n,r}^{(k)}(0)/(L_n^\alpha)^{(k)}(0)$  are uniformly bounded. It is clear that taking derivatives  $k$  times in (7) the connexion coefficients do not change. Then, it could be thought about the possibility to obtain this uniform bound from this formula.

But again we come across the same problem, each term of  $\sum_{i=0}^{r+1} a_{n,r}^i \frac{(L_{n-i}^{\alpha+r+1})^{(k)}(0)}{(L_n^\alpha)^{(k)}(0)}$ , tends to infinity with order  $n^{r+1}$ , but with an alternating sign. To solve this problem, taking



into account the expression relating  $Q_{n,r}^{(k+1)}(0)/(L_n^\alpha)^{(k+1)}(0)$  and  $Q_{n,r}^{(k)}(0)/(L_n^\alpha)^{(k)}(0)$ , (see [5, Lemma 3]), the necessary uniform bound for the ratios could be derived (see [5, Lemma 4]). Then we have

**Theorem 3** *Let  $\{Q_{n,r}\}_{n \geq 0}$  be the sequence of polynomials orthogonal with respect to the inner product (6) with  $(-1)^n/n!$  as leading coefficient. Then,*

$$\lim_n n^{-\alpha} Q_{n,r} \left( \frac{x}{n} \right) = (-1)^{r+1} x^{-\alpha/2} J_{\alpha+2r+2}(2\sqrt{x}),$$

*uniformly on compact subsets of  $\mathbb{C}$ .*

This result gives a positive answer to the conjecture posed in [8]. We would like to note that the approach is totally new and the techniques used in [5] to prove the above Theorem are not a simple generalization of the ones used in [8].

Next, we will show a remarkable difference between the zeros of  $L_n^\alpha$  and the ones of  $Q_{n,r}$  concerning the convergence acceleration to 0. First, we recall (see [23]) that the zeros of the Laguerre polynomials are real, simple and they are located in  $(0, \infty)$ . Denote by  $(x_{n,k})_{k=1}^n$  the zeros of  $L_n^\alpha$  in an increasing order, they satisfy the interlacing property  $0 < x_{n+1,1} < x_{n,1} < x_{n+1,2} < \dots$ , and  $x_{n,k} \xrightarrow[n]{} 0$  for each fixed  $k$ .

Let  $(j_{\alpha,k})_{k \geq 1}$  be the positive zeros of the Bessel function  $J_\alpha$  writing in an increasing order. Then, formula (9) and Hurwitz's theorem lead us to  $nx_{n,k} \xrightarrow[n]{} j_{\alpha,k}$ ,  $k \geq 1$ , and therefore  $x_{n,k} \sim \frac{C_k}{n}$ ,  $k \geq 1$ , where  $C_k$  is a positive constant depending on  $k$ .

Concerning the zeros of  $Q_{n,r}$ , standard arguments (see for instance [10]) allow to establish that  $Q_{n,r}$  has at least  $n - (r + 1)$  zeros with odd multiplicity in  $(0, +\infty)$ , or equivalently  $n - (r + 1)$  changes of sign. Moreover, since  $M_0 > 0$  and the mass point in the discrete part of the inner product belongs to the boundary of  $(0, +\infty)$  then the number of zeros with odd multiplicity is at least  $n - r$  (see [2]).

From Theorem 3 and Hurwitz's theorem, taking into account the multiplicity of 0 as a zero of the limit function in Theorem 3, we achieve

**Corollary 1** *Let  $(\xi_{n,k}^r)_{k=1}^n$  be the zeros of  $Q_{n,r}$ . Then*

$$n \xi_{n,k}^r \xrightarrow[n]{} 0, \quad 1 \leq k \leq r + 1,$$

$$n \xi_{n,k}^r \xrightarrow[n]{} j_{\alpha+2r+2, k-r-1}, \quad k \geq r + 2.$$

**Remark 2.** The presence of the positive masses  $M_i$ ,  $i = 0, \dots, r$ , in the inner product produces a convergence acceleration to 0 of  $r + 1$  zeros of the polynomials  $Q_{n,r}$ .

### 3 Laguerre–Sobolev inner products with holes

Until now, we have assumed that all the masses  $M_i$  in the discrete part of the Sobolev inner product are positive. The possibility of some  $M_i = 0$  has been also dealt in the literature. For instance, the case  $M_0 = 0, M_1 > 0$  ([8]) and similar situations in the non-diagonal case ([7] and [11]) have been analyzed. Very recently, in [12], the authors study the particular case  $M_i = 0, i = 0, \dots, r - 1$ , for the Laguerre–Sobolev type polynomials. The results obtained in all these papers have been generalized in [5], where such a kind of inner products have been called Sobolev inner products with *holes*.

More concretely, we consider the inner product

$$(f, g)_{r,s} = \frac{1}{\Gamma(\alpha + 1)} \int_0^\infty f(x)g(x)x^\alpha e^{-x} dx + \sum_{i=0}^r M_i f^{(i)}(0)g^{(i)}(0) + M_s f^{(s)}(0)g^{(s)}(0), \quad (10)$$

where  $s \geq r + 2$  and  $M_i > 0$  for  $i = 0, \dots, r$  and  $i = s$ .

Observe that we are concerned with inner products of the form

$$(p, q)_{r,s} = (p, q)_r + M_s p^{(s)}(0)q^{(s)}(0), \quad s \geq r + 2,$$

where  $M_s > 0$ , and in  $(\cdot, \cdot)_r$  all the masses are positive. That is, roughly speaking, there is a “hole” in the discrete part of the inner product  $(\cdot, \cdot)_{r,s}$ . We denote by  $\{T_{n,r,s}\}_{n \geq 0}$  the sequence of polynomials orthogonal with respect to the inner product  $(\cdot, \cdot)_{r,s}$  with leading coefficients  $(-1)^n/n!$ .

For this situation, the relative asymptotics and the Mehler-Heine type formulas have been established in [5]. We want to remark that this case has qualitative differences with respect to the case without holes. For example, concerning the convergence acceleration to 0 of the zeros of the polynomials, as we will below.

Arguing as in Lemma 1 it can be proved

**Lemma 2** *Let  $\{T_{n,r,s}\}_{n \geq 0}$  be the sequence of polynomials orthogonal with respect to the inner product (10) with  $(-1)^n/n!$  as leading coefficient. Then the following statements hold:*

(a) *For either  $0 \leq k \leq r$  or  $k = s$ ,*

$$T_{n,r,s}^{(k)}(0) \sim \frac{C_{r,s,k}}{n^{\alpha+2k+1}} (L_n^\alpha)^{(k)}(0),$$

*where  $C_{r,s,k}$  is a nonzero real number independent of  $n$ .*

(b) *For  $k \geq r + 1$  and  $k \neq s$*

$$T_{n,r,s}^{(k)}(0) \sim \frac{k!}{(k - (r + 1))!} \frac{k - s}{\alpha + s + k + 1} \frac{\Gamma(\alpha + k + 1)}{\Gamma(\alpha + r + k + 2)} (L_n^\alpha)^{(k)}(0).$$

(c)

$$(T_{n,r,s}T_{n,r,s})_{r,s} \sim \|L_n^\alpha\|^2.$$

Observe that, as in the complete case (without holes), the addition of the discrete part of the inner product modifies the size of the derivative of order  $k$  only when the corresponding mass  $M_k$  is positive.

Using this lemma the relative asymptotics for these orthogonal polynomials can be deduced:

**Theorem 4** *Let  $\{T_{n,r,s}\}_{n \geq 0}$  be the sequence of o.p. with respect to the inner product (10) with  $(-1)^n/n!$  as leading coefficient . Then*

$$\frac{T_{n,r,s}(x)}{L_n^\alpha(x)} \rightrightarrows 1, \quad x \in \mathbb{C} \setminus [0, \infty).$$

The Mehler–Heine type formula adopts the form

**Theorem 5** *Let  $\{T_{n,r,s}\}_{n \geq 0}$  be the sequence of polynomials orthogonal with respect to the inner product (10) with  $(-1)^n/n!$  as leading coefficient . Then,*

$$\begin{aligned} n^{-\alpha} T_{n,r,s} \left( \frac{x}{n} \right) &\rightrightarrows (-1)^{r+1} x^{-\alpha/2} \\ &\times \left[ \frac{-(s - (r + 1))}{\alpha + r + s + 2} J_{\alpha+2r+2}(2\sqrt{x}) + \sum_{l=2}^{s-r+1} \lambda_l J_{\alpha+2r+2l}(2\sqrt{x}) \right], \quad x \in \mathbb{C}, \end{aligned} \quad (11)$$

where  $\lambda_i$  are nonzero real numbers.

For the particular case  $s = r + 2$ , i.e., when there is a hole of “length one”, the above result generalizes the one obtained in [8]. Theorem 5 also generalizes the corresponding result in [12].

Now, we comment the acceleration of the convergence towards the origin of the zeros of the polynomials  $T_{n,r,s}$ . The quasi-orthogonality of order  $s + 1$  of the sequence  $\{T_{n,r,s}\}_{n \geq 0}$  with respect to the positive weight  $x^{\alpha+s+1}e^{-x}$  assures that  $T_{n,r,s}$  has at least  $n - (s + 1)$  changes of sign in  $(0, +\infty)$ . However, in [2] the authors proved that the number of zeros in  $(0, +\infty)$  does not depend on the order of the derivatives but on the number of terms in the discrete part of the inner product. So,  $T_{n,r,s}$  has at least  $n - (r + 1)$  zeros with odd multiplicity in  $(0, +\infty)$ . Proceeding as in Corollary 1, we get:

**Corollary 2** *Let  $(\zeta_{n,k}^{r,s})_{k=1}^n$  be the zeros of  $T_{n,r,s}$ . Then*

$$\begin{aligned} n \zeta_{n,k}^{r,s} &\rightarrow 0, \quad 1 \leq k \leq r + 1, \\ n \zeta_{n,k}^{r,s} &\rightarrow j_{\alpha+2r+2,k-r-1}, \quad k \geq r + 2. \end{aligned}$$

**Remark 3.** We want to highlight that this result is in a way surprising since it does not depend on the number of terms in the discrete part, but on the position of the hole. So, despite the presence of the mass  $M_s$ , there only exists an acceleration of the convergence of  $r + 1$  zeros such as it occurs in the case of the inner products without holes. That is, the convergence acceleration to 0 of the zeros of the polynomials  $Q_{n,r}$  and  $T_{n,r,s}$  is the same and the addition of a mass  $M_s$  *after a hole* in the inner product does not affect the convergence acceleration to 0.

#### 4 Generalized Hermite–Sobolev type polynomials

As a consequence of the previous results, asymptotic properties for the orthogonal polynomials  $S_{n,r}^\mu$  associated with the inner product

$$(p, q) = \int_{\mathbb{R}} p(x)q(x)|x|^{2\mu} e^{-x^2} dx + \sum_{i=0}^{2r+1} M_i p^{(i)}(0) q^{(i)}(0), \quad (12)$$

with  $\mu > -1/2$  and  $M_i > 0$ ,  $i = 0, \dots, 2r + 1$ , can be established. We assume that the leading coefficient of  $S_{n,r}^\mu$  is  $2^n$ .

Remind that the polynomials  $H_n^\mu$  orthogonal with respect to the weight  $|x|^{2\mu} e^{-x^2}$  ( $\mu > -1/2$ ) are called *generalized Hermite polynomials*. So, we are concerned with generalized Hermite–Sobolev type orthogonal polynomials.

Notice that in this case the polynomials  $S_{n,r}^\mu$  are symmetric, that is,  $S_{n,r}^\mu(-x) = (-1)^n S_{n,r}^\mu(x)$ , and because of this symmetry, we can transform the inner product (12) into an inner product like (6) and so we can establish a simple relation between the polynomials  $S_{n,r}^\mu$  and the polynomials  $Q_{n,r}$  considered before. This technique is known as a symmetrization process. In fact, in [10] this process is considered for standard inner products associated with positive measures. The simplest case of this situation is the relation between Laguerre polynomials and Hermite polynomials, that is (see [10] or [23]), for  $n \geq 0$ ,

$$H_{2n}(x) = (-1)^n 2^{2n} n! L_n^{-1/2}(x^2), \quad H_{2n+1}(x) = (-1)^n 2^{2n+1} n! x L_n^{1/2}(x^2).$$

Later in [3] the authors generalize the symmetrization process in the framework of Sobolev type orthogonal polynomials, (see Theorem 2 in [3]). Thus,

$$S_{2n,r}^\mu(x) = (-1)^n 2^{2n} n! Q_{n,r}^{\mu-1/2}(x^2), \quad S_{2n+1,r}^\mu(x) = (-1)^n 2^{2n+1} n! x Q_{n,r}^{\mu+1/2}(x^2)$$

where  $\{Q_{n,r}^{\mu-1/2}\}_{n \geq 0}$  (respectively,  $\{Q_{n,r}^{\mu+1/2}\}_{n \geq 0}$ ) is the sequence of polynomials orthogonal with respect to an inner product like (6) with  $\alpha = \mu - 1/2$  (respectively,  $\alpha = \mu + 1/2$ ) and leading coefficient  $(-1)^n/n!$ .

Using this symmetrization process, the relative asymptotics and the Mehler–Heine type formulas for generalized Hermite–Sobolev type polynomials can be proved.

**Proposition 2** Let  $\{S_{n,r}^\mu\}_{n \geq 0}$  be the sequence of polynomials orthogonal with respect to the inner product (12) with  $2^n$  as leading coefficient. Then,

(a)

$$\frac{S_{n,r}^\mu(x)}{H_n^\mu(x)} \Rightarrow 1, \quad x \in \mathbb{C} \setminus \mathbb{R}.$$

(b)

$$n^{-\mu+1/2} S_{2n,r}^\mu \left( \frac{x}{2\sqrt{n}} \right) \Rightarrow (-1)^{r+1} \left( \frac{x}{2} \right)^{-\mu+1/2} J_{\mu+2r+3/2}(x), \quad x \in \mathbb{C}$$

$$n^{-\mu+1/2} S_{2n+1,r}^\mu \left( \frac{x}{2\sqrt{n}} \right) \Rightarrow (-1)^{r+1} \left( \frac{x}{2} \right)^{-\mu+1/2} J_{\mu+2r+5/2}(x), \quad x \in \mathbb{C}.$$

**Remark 4.** These results generalize some of the results in [4] and solve the conjecture posed there.

Using a symmetrization process, relative asymptotics and Mehler–Heine type formulas for generalized Hermite–Sobolev polynomials with holes in the discrete part of the inner product can be deduced.

Finally, we hope this method can be used with other measures with unbounded support for which we have quite less explicit information about the corresponding orthogonal polynomials.

## Acknowledgements

This work has been partially supported by MICINN of Spain under Grant MTM2009-12740-C03-03, FEDER funds (EU), and the Diputación General de Aragón, project E-64.

## References

- [1] M. Alfaro, M. Álvarez de Morales and M.L. Rezola, *Orthogonality of the Jacobi polynomials with negative parameters*, J. Comput. Appl. Math. **145** (2002), 379–386.
- [2] M. Alfaro, G. López and M.L. Rezola, *Some properties of zeros of Sobolev-type orthogonal polynomials*, J. Comput. Appl. Math. **69** (1996), 171–179.
- [3] M. Alfaro, F. Marcellán, H.G. Meijer and M.L. Rezola, *Symmetric orthogonal polynomials for Sobolev-type inner products*, J. Math. Anal. Appl. **184** (1994), 360–381.
- [4] M. Alfaro, J.J. Moreno–Balcázar, A. Peña and M.L. Rezola, *Asymptotics for a generalization of Hermite polynomials*. Asymptotic Anal. **66** (2010), 103–117.
- [5] M. Alfaro, J.J. Moreno–Balcázar, A. Peña and M.L. Rezola, *A new approach to the asymptotics for Sobolev orthogonal polynomials*. Submitted for publication, 2010. arXiv 1003.3336v1

- [6] R. Álvarez–Nodarse and F. Marcellán, *A generalization of the classical Laguerre polynomials*, Rend.Circ. Mat. Palermo (2) **44** (1995), 315–329.
- [7] R. Álvarez–Nodarse and F. Marcellán, *A generalization of the classical Laguerre polynomials: asymptotic properties and zeros*, Appl. Anal. **62** (1996), 349–366.
- [8] R. Álvarez–Nodarse and J.J. Moreno–Balcázar, *Asymptotic properties of generalized Laguerre orthogonal polynomials*, Indag. Mathem., N.S. **15** (2004), 151–165.
- [9] K.E. Atkinson and O. Hansen, *Solving the nonlinear Poisson equation on the unit disk*, J. Integral Eq. and Appl., **17** (2006), 223–241.
- [10] T.S. Chihara, *An Introduction to Orthogonal Polynomials*. Gordon & Breach, New York, 1978.
- [11] H. Dueñas and F. Marcellán, *Asymptotic behaviour of the Laguerre–Sobolev–type orthogonal polynomials. A nondiagonal case*, J. Comput. Appl. Math. (2009), doi:10.1016/j.cam.2009.07.055.
- [12] H. Dueñas and F. Marcellán, *The Laguerre–Sobolev–Type Orthogonal Polynomials*, J. Approx. Theory **162** (2010), 421–440.
- [13] D. Evans, L.L. Littlejohn, F. Marcellán, C. Market and A. Ronveaux *On recurrence relations for Sobolev polynomials*, SIAM J. Math. Anal. **26** (1995), 446–467.
- [14] L. Fernández, F. Marcellán, T.E. Pérez and M.A. Piñar *Recent trends on two variable orthogonal polynomials*, Contemp. Math. **509** (2010), 59–86.
- [15] R. Koekoek, *Generalizations of Laguerre polynomials*, J. Math. Anal. Appl. **153** (1990), 576–590.
- [16] R. Koekoek and H.G. Meijer, *A generalization of Laguerre polynomials*, SIAM J. Math. Anal. **24** (1993), 768–782.
- [17] K.H. Kwon and L.L. Littlejohn, *The orthogonality of the Laguerre polynomials  $\{L_n^{(-k)}(x)\}$  for a positive integer  $k$* , Ann. Numer. Math. **2** (1995), 289–304.
- [18] K.H. Kwon, L.L. Littlejohn and G.J. Yoon, *Ghost matrices and a characterization of symmetric Sobolev bilinears forms*, Linear Algebra Appl. **431** (2009), 104–119.
- [19] J.K. Lee and L. L. Littlejohn *Sobolev orthogonal polynomials in two variables and second order partial differential equations*, J. Math. Anal. Appl. **322** (2006), 1001–1017.
- [20] G. López, F. Marcellán and W. Van Assche, *Relative asymptotics for polynomials orthogonal with respect to a discrete Sobolev inner product*, Constr. Approx. **11** (1995), 107–137.

- [21] F. Marcellán and J.J. Moreno–Balcázar, *Asymptotics and zeros of Sobolev orthogonal polynomials on unbounded supports*, Acta Appl. Math. **94** (2006), 163–192.
- [22] I.A. Rocha, F. Marcellán and L. Salto, *Relative asymptotics and Fourier series of orthogonal polynomials with a discrete Sobolev inner product*, J. Approx. Theory **121** (2003), 336–356.
- [23] G. Szegő, *Orthogonal Polynomials*, Amer. Math. Soc. Colloq. Publ. vol. **23**, Amer. Math. Soc., Providence R.I., 1975. Fourth Edition.
- [24] Y. Xu, *A family of Sobolev orthogonal polynomials on the unit ball*, J. Approx. Theory, **138** (2006), 232–241.

